Exercise 8 from WF122

Our next exercise we have used before:

```
(0) X \Rightarrow \mathbf{true}.
```

Since the only given properties of **true** in WF122 relate it to \equiv , $\not\equiv$, \lor , and \land , we should rewrite (0) by unfolding \Rightarrow in terms of these. This yields the following attempts:

```
X \Rightarrow \mathbf{true}
\equiv \{ \Rightarrow \text{ into } \neg / \lor \} \}
\neg X \lor \mathbf{true}
\equiv \{ \text{ predicate calculus } \} \}
\mathbf{true}
\equiv \{ \Rightarrow \text{ into } \equiv / \lor \} \}
\mathbf{true} \equiv X \lor \mathbf{true}
\equiv \{ \text{ predicate calculus } \} \}
\mathbf{true} \equiv \mathbf{true}
\equiv \{ \text{ predicate calculus } \} \}
```

and:

true

EX8-1

$$X \Rightarrow \mathbf{true}$$

$$\equiv \{ \Rightarrow \text{ into } \equiv / \land \} \}$$

$$X \equiv X \land \mathbf{true}$$

$$\equiv \{ \text{ predicate calculus } \}$$

$$X \equiv X$$

$$\equiv \{ \text{ predicate calculus } \}$$

$$\mathbf{true} \qquad .$$

That was an exercise in just writing down "the only thing I could do" .

NYC, 5 November 2009

Jeremy Weissmann jeremy @ mathmeth.com